

MEAN CURVATURE FLOW OF HIGHER CODIMENSION IN HYPERBOLIC SPACES

KEFENG LIU, HONGWEI XU, FEI YE, AND ENTAO ZHAO

ABSTRACT. In this paper we investigate the convergence for the mean curvature flow of closed submanifolds with arbitrary codimension in space forms. Particularly, we prove that the mean curvature flow deforms a closed submanifold satisfying a pinching condition in a hyperbolic space form to a round point in finite time.

1. INTRODUCTION

In this paper, we study the convergence of the mean curvature flow of submanifolds in space forms. Let $F : M^n \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth immersion from an n -dimensional closed Riemannian manifold M^n to an $(n+d)$ -dimensional complete simply connected space form $\mathbb{F}^{n+d}(c)$ with constant sectional curvature c . Consider a one-parameter family of smooth immersions $F : M \times [0, T) \rightarrow \mathbb{F}^{n+d}(c)$ satisfying

$$(1.1) \quad \begin{cases} \frac{\partial}{\partial t} F(x, t) &= H(x, t), \\ F(x, 0) &= F(x), \end{cases}$$

where $H(x, t)$ is the mean curvature vector of $F_t(M)$ and $F_t(x) = F(x, t)$. We call $F : M \times [0, T) \rightarrow \mathbb{F}^{n+d}(c)$ the mean curvature flow with initial value F .

The mean curvature flow was proposed by Mullins [17] to describe the formation of grain boundaries in annealing metals. In [3], Brakke introduced the motion of a submanifold by its mean curvature in arbitrary codimension and constructed a generalized varifold solution for all time. For the classical solution of the mean curvature flow, most works have been done on hypersurfaces. Huisken [11, 12] showed that if the initial hypersurface in a Riemannian manifold is uniformly convex, then the mean curvature flow converges to a round point in finite time. Later, Huisken [13] extend this result to hypersurfaces satisfying a pinching condition in a sphere. Many other beautiful results have been obtained, and there are various approaches to study the mean curvature flow of hypersurfaces (see [6, 7], etc.). For the mean curvature flow of submanifolds in higher codimension, some special cases have been studied, see [19, 20, 21, 22, 23, 24] etc. for example. Recently, Andrews-Baker [1] proved a convergence theorem for the mean curvature flow of closed submanifolds satisfying a pinching condition in the Euclidean space. In [2], Baker proved a convergence result for the mean curvature flow of submanifolds in a sphere. In this paper, we study the mean curvature flow of closed submanifolds in hyperbolic

2000 *Mathematics Subject Classification.* 53C44, 53C40.

Key words and phrases. Mean curvature flow; higher codimension; submanifolds; convergence; second fundamental form.

Research supported by the National Natural Science Foundation of China, Grant No. 11071211; the Trans-Century Training Programme Foundation for Talents by the Ministry of Education of China, and the China Postdoctoral Science Foundation, Grant No. 20090461379.

spaces and extend the convergence result in [1, 2] to the mean curvature flow of arbitrary codimension in space forms.

Theorem 1.1. *Let $F : M^n \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth closed submanifold in a hyperbolic space with constant curvature $c < 0$. Assume F satisfies*

$$(1.2) \quad |A|^2 \leq \begin{cases} \frac{4}{3n}|H|^2 + \frac{n}{2}c, & n = 2, 3, \\ \frac{1}{n-1}|H|^2 + 2c, & n \geq 4. \end{cases}$$

Then the mean curvature flow with F as initial value converges to a round point in finite time.

As an immediate consequence of Theorem 1.1, we obtain the following differentiable sphere theorem.

Corollary 1.2. *Let $F : M^n \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth closed submanifold in a hyperbolic space with constant curvature $c < 0$. Assume F satisfies*

$$|A|^2 \leq \begin{cases} \frac{4}{3n}|H|^2 + \frac{n}{2}c, & n = 2, 3, \\ \frac{1}{n-1}|H|^2 + 2c, & n \geq 4. \end{cases}$$

Then M is diffeomorphic to the unit n -sphere.

Remark 1.3. This differentiable sphere theorem was also obtained by Gu and Xu [10, 25] provided the submanifold is simply connected. In fact, they proved the sphere theorem for submanifolds in a Riemannian manifold by using a different method. For more sphere theorems of submanifolds, we refer the readers to [1, 2, 9, 10, 16, 18, 25, 28], etc.

Combining Theorem 1.1 and the convergence results in [1, 2], we obtain the following theorem.

Theorem 1.4. *Let $F : M^n \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth closed submanifold in a complete simply connected space form with $|H|^2 + n^2c > 0$. Assume F satisfies*

$$(1.3) \quad |A|^2 \leq \begin{cases} \frac{4}{3n}|H|^2 + \frac{1}{12}[7n - 4 + \operatorname{sgn}(c)(n - 4)]c, & n = 2, 3, \\ \frac{1}{n-1}|H|^2 + 2c, & n \geq 4. \end{cases}$$

Then either $F_t(M)$ converges to a round point in finite time, or $c > 0$ and $F_t(M)$ converges to a total geodesic sphere in $\mathbb{F}^{n+d}(c)$ as $t \rightarrow \infty$.

Remark 1.5. For $c > 0$, $|H|^2 + n^2c > 0$ is automatically satisfied. For $c = 0$, $|H|^2 + n^2c > 0$ is equivalent to that the mean curvature is nowhere vanishing. For $c < 0$, $|H|^2 + n^2c > 0$ is implied by condition (1.3).

Remark 1.6. For $c > 0$, the maximal existence time of the mean curvature flow may be finite or infinite. For $c \leq 0$, the mean curvature flow with a closed initial submanifold always has finite maximal existence time.

2. BASIC EQUATIONS

Let $F : M \times [0, T) \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth mean curvature flow with initial closed immersion $F_0 : M \rightarrow \mathbb{F}^{n+d}(c)$. Denote by $g(t)$ and $d\mu_t$ the induced metric and the volume form on M . Let A and H be the second fundamental form and

the mean curvature vector of M in $\mathbb{F}^{n+d}(c)$, respectively. We shall make use of the following convention on the range of indices.

$$1 \leq i, j, k, \dots \leq n, \quad 1 \leq A, B, C, \dots \leq n+d \quad \text{and} \quad n+1 \leq \alpha, \beta, \gamma, \dots \leq n+d.$$

As in [1, 3], we consider the evolution on the spatial tangent bundle. Choose a local orthonormal frame $\{e_i\}$ for the spatial tangent bundle and a local orthonormal frame $\{\nu_\alpha\}$ for the normal bundle. Let $\{\omega_i\}$ be the dual frame of $\{e_i\}$. Then A and H can be written as

$$A = \sum_{i,j,\alpha} h_{ij\alpha} \omega_i \otimes \omega_j \otimes \nu_\alpha = \sum_{i,j} h_{ij} \omega_i \otimes \omega_j, \quad H = \sum_{\alpha} H_{\alpha} \nu_{\alpha}.$$

We have the following evolution equations.

$$(2.1) \quad \frac{\partial}{\partial t} |A|^2 = \Delta |A|^2 - 2|\nabla A|^2 + 2R_1 + 4c|H|^2 - 2nc|A|^2,$$

$$(2.2) \quad \frac{\partial}{\partial t} |H|^2 = \Delta |H|^2 - 2|\nabla H|^2 + 2R_2 + 2nc|H|^2,$$

where

$$(2.3) \quad R_1 = \sum_{\alpha,\beta} \left(\sum_{i,j} h_{ij\alpha} h_{ij\beta} \right)^2 + |R^\perp|^2,$$

$$(2.4) \quad |R^\perp|^2 = \sum_{i,j,\alpha,\beta} \left(\sum_p (h_{ip\alpha} h_{jp\beta} - h_{jp\alpha} h_{ip\beta}) \right)^2,$$

$$(2.5) \quad R_2 = \sum_{i,j} \left(\sum_{\alpha} H_{\alpha} h_{ij\alpha} \right)^2.$$

The contracted form of Simons' identity for traceless second fundamental form $\mathring{A} := A - \frac{1}{n}g \otimes H$ is

$$(2.6) \quad \frac{1}{2} \Delta |\mathring{A}|^2 = \mathring{h}_{ij} \nabla_i \nabla_j H + |\nabla \mathring{A}|^2 + Z + nc|\mathring{A}|^2.$$

Here

$$(2.7) \quad Z = -R_1 + \sum_{i,j,p,\alpha,\beta} H_{\alpha} h_{ip\alpha} h_{ij\beta} h_{pj\beta}.$$

We also have the following inequality.

$$(2.8) \quad |\nabla A|^2 \geq \frac{3}{n+2} |\nabla H|^2.$$

3. PRESERVED CURVATURE PINCHING CONDITION

Now we prove that the pinching condition (1.2) is preserved under the mean curvature flow with arbitrary codimension in the hyperbolic space.

Lemma 3.1. *For $c < 0$ and $n \geq 2$, if the initial immersion satisfies (1.2), then this condition is preserved along the mean curvature flow.*

Proof. We consider $Q = |A|^2 - \alpha|H|^2 - \beta c$, where the constants

$$\alpha \leq \begin{cases} \frac{4}{3n}, & n = 2, 3, \\ \frac{1}{n-1}, & n \geq 4, \end{cases} \quad \text{and} \quad \beta \geq \begin{cases} \frac{n}{2}, & n = 2, 3, \\ 2, & n \geq 4. \end{cases}$$

By (2.1) and (2.2) we have

$$(3.1) \quad \begin{aligned} \frac{\partial}{\partial t} Q &= \Delta Q - 2(|\nabla A|^2 - \alpha|\nabla H|^2) \\ &\quad + 2R_1 - 2\alpha R_2 - 2nc|\mathring{A}|^2 - 2n\left(\alpha - \frac{1}{n}\right)c|H|^2. \end{aligned}$$

We only have to show that if $Q = 0$ at a point $x \in M$, then

$$2R_1 - 2\alpha R_2 - 2nc|\mathring{A}|^2 - 2n\left(\alpha - \frac{1}{n}\right)c|H|^2 \leq 0$$

holds at x . We also have $H \neq 0$ at x . Choose $\{\nu_\alpha\}$ such that $\nu_{n+1} = \frac{H}{|H|}$. Let $A_H = \sum_{i,j} h_{ij,n+1} \omega_i \otimes \omega_j$. Set $\mathring{A}_H = A_H - \frac{|H|}{n} \text{Id}$ and $|\mathring{A}_H|^2 = |\mathring{A}|^2 - |\mathring{A}_H|^2$.

We replace $|H|^2$ with $\frac{|\mathring{A}|^2 - \beta c}{\alpha - \frac{1}{n}}$. Then

$$(3.2) \quad \begin{aligned} &2R_1 - 2\alpha R_2 - 2nc|\mathring{A}|^2 - 2n\left(\alpha - \frac{1}{n}\right)c|H|^2 \\ &\leq 2|\mathring{A}_H|^2 - 2\left(\alpha - \frac{1}{n}\right)|\mathring{A}_H|^2|H|^2 + \frac{2}{n}|\mathring{A}_H|^2|H|^2 - \frac{2}{n}\left(\alpha - \frac{1}{n}\right)|H|^4 \\ &\quad + 8|\mathring{A}_H|^2|\mathring{A}_I|^2 + 3|\mathring{A}_I|^2 - 2nc(|\mathring{A}_H|^2 + |\mathring{A}_I|^2) - 2n\left(\alpha - \frac{1}{n}\right)c|H|^2 \\ &= \left(6 - \frac{2}{n(\alpha - \frac{1}{n})}\right)|\mathring{A}_H|^2|\mathring{A}_I|^2 + \left(3 - \frac{2}{n(\alpha - \frac{1}{n})}\right)|\mathring{A}_I|^4 \\ &\quad + \left(2\beta - 4n + \frac{2\beta}{n(\alpha - \frac{1}{n})}\right)c|\mathring{A}_H|^2 + 4\left(\frac{\beta}{n(\alpha - \frac{1}{n})} - n\right)c|\mathring{A}_I|^2 \\ &\quad - 2\beta\left(\frac{\beta}{n(\alpha - \frac{1}{n})} - n\right)c^2. \end{aligned}$$

By the definition of α and β , we know that the right hand side of (3.2) is nonpositive for $n \geq 2$. This completes the proof of the lemma. \square

For $\epsilon > 0$, set

$$\alpha_\epsilon = \begin{cases} \frac{4}{3n+n\epsilon}, & n = 2, 3, \\ \frac{1}{n-1+\epsilon}, & n \geq 4, \end{cases} \quad \text{and} \quad \beta_\epsilon = \begin{cases} \frac{n}{2}(1+\epsilon), & n = 2, 3, \\ 2(1+\epsilon), & n \geq 4. \end{cases}$$

If the initial immersion satisfies $|A|^2 < \frac{4}{3n}|H|^2 + \frac{n}{2}c$ for $n = 2, 3$, and $|A|^2 < \frac{1}{n-1}|H|^2 + 2c$ for $n \geq 4$, then there exists an $\epsilon > 0$ such that $|A|^2 \leq \alpha_\epsilon|H|^2 + \beta_\epsilon c$ holds on M_0 . From the proof of Lemma 3.1, this inequality also holds for $t > 0$. On the other hand, if $|A|^2 = \frac{4}{3n}|H|^2 + \frac{n}{2}c$ for $n = 2, 3$, or $|A|^2 = \frac{1}{n-1}|H|^2 + 2c$ for $n \geq 4$ holds somewhere on M_0 , then by the maximum principle, we see that either the equality holds everywhere on M_0 , or the strict inequality holds everywhere for $t > 0$. For the first case, we have $\nabla A = 0$ and $\mathring{A}_I = 0$ on M_0 . By [8], M_0 lies in an $(n+1)$ -dimensional total geodesic submanifold of $\mathbb{F}^{n+d}(c)$. Since $\nabla A = 0$, from Theorem 4 of [15], M_0 is either locally isometric to an Euclidean space, or

locally isometric to a product $\mathbb{F}^k(c_1) \times \mathbb{F}^{n-k}(c_2)$ for some $c_1 > 0$, $c_2 < 0$ and $k = 0, \dots, n$. Since M_0 is closed, we see that M_0 is a totally umbilical sphere. Then $|A|^2 \leq \alpha_\epsilon |H|^2 + \beta_\epsilon c$ holds on M_0 for some $\epsilon > 0$. For the second case, we see that after a short time, we also have $|A|^2 \leq \alpha_\epsilon |H|^2 + \beta_\epsilon c$ for some $\epsilon > 0$. Hence, we may assume that $|A|^2 \leq \alpha_\epsilon |H|^2 + \beta_\epsilon c$ for some $\epsilon \in (0, 1)$ and $t \geq t_0 > 0$.

4. PINCHING OF \mathring{A} ALONG THE MEAN CURVATURE FLOW

Assume that $c < 0$. We prove a pinching estimate for the traceless second fundamental form, which guarantees that M_t becomes spherical along the mean curvature flow.

Theorem 4.1. *There are positive constants C_0 and σ_0 independent of t such that*

$$(4.1) \quad |\mathring{A}|^2 \leq C_0 |H|^{2-\sigma_0}$$

holds along the mean curvature flow.

Proof. We consider the function $f_\sigma = \frac{|\mathring{A}|^2}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}}$, where $\sigma \in (0, 1)$ and

$$a = \begin{cases} \frac{1}{3n+n\epsilon}, & n = 2, 3 \\ \frac{1}{n(n-1+\epsilon)}, & n \geq 4. \end{cases}$$

Notice that

$$(4.2) \quad \begin{aligned} & a|H|^2 + \beta_\epsilon c - \left(\left(\alpha_\epsilon - \frac{1}{n} \right) |H|^2 + \beta_\epsilon c \right) \\ & \geq \frac{\epsilon}{3n+n\epsilon} |H|^2 \\ & \geq \frac{\epsilon}{3n+n\epsilon} \cdot \beta(-c) \\ & > 0 \end{aligned}$$

for $n = 2, 3$, and

$$(4.3) \quad \begin{aligned} & a|H|^2 + \beta_\epsilon c - \left(\left(\alpha_\epsilon - \frac{1}{n} \right) |H|^2 + \beta_\epsilon c \right) \\ & \geq \frac{\epsilon}{n(n-1+\epsilon)} |H|^2 \\ & \geq \frac{\epsilon}{n(n-1+\epsilon)} \cdot \beta(-c) \\ & > 0 \end{aligned}$$

for $n \geq 4$. So f_σ is well-defined. From (4.2) and (4.3) we also have

$$(4.4) \quad a|H|^2 + \beta_\epsilon c \geq b|H|^2,$$

where

$$b = \begin{cases} \frac{\epsilon}{3n+n\epsilon}, & n = 2, 3 \\ \frac{\epsilon}{n(n-1+\epsilon)}, & n \geq 4. \end{cases}$$

By a similar computation as in [1], we have

$$\begin{aligned}
(4.5) \quad \frac{\partial}{\partial t} f_\sigma &= \Delta f_\sigma + \frac{2a(1-\sigma)}{a|H|^2 + \beta_\epsilon c} \langle \nabla |H|^2, \nabla f_\sigma \rangle \\
&\quad - \frac{2}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} \left(|\nabla A|^2 - \frac{1}{n} |\nabla H|^2 - \frac{a|\mathring{A}|^2}{a|H|^2 + \beta_\epsilon c} |\nabla H|^2 \right) \\
&\quad - \frac{4a^2\sigma(1-\sigma)}{(a|H|^2 + \beta_\epsilon c)^2} f_\sigma |H|^2 \cdot |\nabla |H||^2 - \frac{2a\sigma f_\sigma}{a|H|^2 + \beta_\epsilon c} |\nabla H|^2 \\
&\quad + \frac{2}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} \left(R_1 - \frac{1}{n} R_2 - \frac{aR_2|\mathring{A}|^2}{a|H|^2 + \beta_\epsilon c} - nc|\mathring{A}|^2 \right. \\
&\quad \left. - \frac{an(1-\sigma)c|\mathring{A}|^2|H|^2}{a|H|^2 + \beta_\epsilon c} \right) \\
&\quad + \frac{2a\sigma R_2 f_\sigma}{a|H|^2 + \beta_\epsilon c}.
\end{aligned}$$

By (2.8), we have

$$\begin{aligned}
(4.6) \quad & |\nabla A|^2 - \frac{1}{n} |\nabla H|^2 - \frac{a|\mathring{A}|^2}{a|H|^2 + \beta_\epsilon c} |\nabla H|^2 \\
& \geq \left(\frac{3}{n+2} - \frac{1}{n} - \frac{a\left((\alpha_\epsilon - \frac{1}{n})H^2 + \beta_\epsilon c\right)}{a|H|^2 + \beta_\epsilon c} \right) |\nabla H|^2 \\
& \geq \left(\frac{3}{n+2} - \frac{1}{n} - a \right) |\nabla H|^2 \\
& := \epsilon_\nabla |\nabla H|^2.
\end{aligned}$$

Here ϵ_∇ is a positive constant for $n \geq 2$.

We also have the following estimate.

$$\begin{aligned}
(4.7) \quad & R_1 - \frac{1}{n} R_2 - \frac{aR_2|\mathring{A}|^2}{a|H|^2 + \beta_\epsilon c} \\
& \leq R_1 - \frac{1}{n} R_2 - \frac{R_2|\mathring{A}|^2}{|H|^2} \\
& = R_1 - \frac{R_2|A|^2}{|H|^2} \\
& \leq |\mathring{A}_H|^4 - 2 \left(\frac{|A|^2}{|H|^2} - \frac{2}{n} \right) |\mathring{A}_H|^2 |H|^2 - \frac{2}{n} \left(\frac{|A|^2}{|H|^2} - \frac{1}{n} \right) |H|^4 \\
& \quad - 4|\mathring{A}_H|^2 |\mathring{A}_I|^2 - \frac{3}{2} |\mathring{A}_I|^4 \\
& \leq 0.
\end{aligned}$$

In (4.7) we have used the pinching condition $|A|^2 \leq \alpha_\epsilon |H|^2 + \beta_\epsilon c < \alpha_\epsilon |H|^2$ for $\epsilon \in (0, 1)$.

By (4.4), we have

$$\begin{aligned}
 & -nc|\mathring{A}|^2 - \frac{an(1-\sigma)c|\mathring{A}|^2|H|^2}{a|H|^2 + \beta_\epsilon c} \\
 (4.8) \quad & \leq -nc|\mathring{A}|^2 - \frac{an(1-\sigma)c|\mathring{A}|^2(a|H|^2 + \beta_\epsilon c)}{b(a|H|^2 + \beta_\epsilon c)} \\
 & \leq -nc|\mathring{A}|^2 - \frac{anc}{b}|\mathring{A}|^2 \\
 & := \bar{b}|\mathring{A}|^2.
 \end{aligned}$$

For the last term of right hand side of (4.5), we have by (4.4)

$$(4.9) \quad \frac{2a\sigma R_2 f_\sigma}{a|H|^2 + \beta_\epsilon c} \leq \frac{2a\sigma|H|^2|A|^2 f_\sigma}{a|H|^2 + \beta_\epsilon c} \leq \frac{2a\sigma}{b}|A|^2 f_\sigma := \tilde{b}\sigma|A|^2 f_\sigma.$$

Combining (4.5), (4.6), (4.7), (4.8) and (4.9), we have

$$\begin{aligned}
 (4.10) \quad \frac{\partial}{\partial t} f_\sigma & \leq \triangle f_\sigma + \frac{2a(1-\sigma)}{a|H|^2 + \beta_\epsilon c} \langle \nabla|H|^2, \nabla f_\sigma \rangle - \frac{2\epsilon_\nabla}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 \\
 & \quad + 2\bar{b}f_\sigma + \tilde{b}\sigma|A|^2 f_\sigma.
 \end{aligned}$$

To deal with the last term of the right hand side of (4.10), we need the following estimate.

Proposition 4.2. *There exists a positive constant ε independent of t such that*

$$(4.11) \quad Z + nc|\mathring{A}|^2 \geq \varepsilon|\mathring{A}|^2(a|H|^2 + \beta_\epsilon c)$$

holds for $t \geq t_0$.

Proof. By the argument in the proof of Lemma 5.4 in [2], we only have to show

$$-\frac{\beta_\epsilon}{\alpha_\epsilon - \frac{1}{n}} \left(\frac{1}{n} - \frac{n-2}{2n(n-1)} \right) + n \leq 0.$$

This is true by our choice of α_ϵ and β_ϵ . □

Proposition 4.3. *For any $\eta > 0$, $p \geq 2$ and $t \geq t_0$, we have*

$$\begin{aligned}
 (4.12) \quad \int_{M_t} f_\sigma^p(a|H|^2 + \beta_\epsilon c) d\mu_t & \leq \frac{2p\eta + 5}{b\varepsilon} \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\
 & \quad + \frac{2(p-1)}{b\eta\varepsilon} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t.
 \end{aligned}$$

Proof. We have the following estimate.

$$\begin{aligned}
& 2 \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} Z d\mu_t + 2nc \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\mathring{A}|^2 d\mu_t \\
& \leq 2(p-1) \int_{M_t} \frac{f_\sigma^{p-2}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla f_\sigma| |\mathring{A}| |\nabla H| d\mu_t \\
& \quad + \frac{2(n-1)}{n} \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\
& \quad + 4 \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{2-\sigma}} |H| |\mathring{A}| |\nabla H|^2 d\mu_t \\
& \quad + 4(1-\sigma)(p-2) \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)} |H| |\nabla H| |\nabla f_\sigma| d\mu_t \\
& \quad + 4 \int_{M_t} \frac{f_\sigma^p}{(a|H|^2 + \beta_\epsilon c)^2} |H|^2 |\nabla H|^2 d\mu_t.
\end{aligned} \tag{4.13}$$

Since $|\mathring{A}|^2 \leq f_\sigma(a|H|^2 + \beta_\epsilon c)^{1-\sigma}$ and $f_\sigma \leq (a|H|^2 + \beta_\epsilon c)^\sigma$, by choosing $\sigma \in (0, 1)$ we have the following estimates.

$$\begin{aligned}
& 2(p-1) \int_{M_t} \frac{f_\sigma^{p-2}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla f_\sigma| |\mathring{A}| |\nabla H| d\mu_t \\
& \leq \frac{1-\sigma}{\eta} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t + (p-1)\eta \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t,
\end{aligned} \tag{4.14}$$

$$\begin{aligned}
& 4 \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{2-\sigma}} |H| |\mathring{A}| |\nabla H|^2 d\mu_t \\
& \leq \frac{4}{b} \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t,
\end{aligned} \tag{4.15}$$

$$\begin{aligned}
& 4(p-2) \int_{M_t} \frac{f_\sigma^{p-1}}{a|H|^2 + \beta_\epsilon c} |H| |\nabla H| |\nabla f_\sigma| d\mu_t \\
& \leq (p-2) \int_{M_t} \frac{1}{a|H|^2 + \beta_\epsilon c} \left(\frac{2}{\eta} f_\sigma^{p-2} |H|^2 |\nabla f_\sigma|^2 + 2\eta f_\sigma^p |\nabla H|^2 \right) d\mu_t \\
& \leq \frac{2(p-2)}{b\eta} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t \\
& \quad + 2(p-2)\eta \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t,
\end{aligned} \tag{4.16}$$

$$4 \int_{M_t} \frac{f_\sigma^p}{(a|H|^2 + \beta_\epsilon c)^2} |H|^2 |\nabla H|^2 d\mu_t \leq \frac{4}{b} \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t. \tag{4.17}$$

In (4.15), (4.16) and (4.17) we have used (4.4).

By (4.11), we have

$$\begin{aligned}
& 2 \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} (Z + 2nc|\mathring{A}|^2) d\mu_t \\
& \geq 2\varepsilon \int_{M_t} f_\sigma^p (a|H|^2 + \beta_\epsilon c) d\mu_t.
\end{aligned} \tag{4.18}$$

Combining (4.13)-(4.18), we obtain

$$(4.19) \quad \begin{aligned} 2\varepsilon \int_{M_t} f_\sigma^p (a|H|^2 + \beta_\epsilon c) d\mu_t &\leq \frac{3p\eta + 10}{b} \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\ &\quad + \frac{3(p-1)}{b\eta} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t. \end{aligned}$$

Dividing through by 2ε completes the proof. \square

Now we show that the L^p -norm of f_σ is bounded for sufficiently high p .

Lemma 4.4. *For any $p \geq \max\{2, \frac{8}{b\epsilon\nabla} + 1\}$ and $\sigma \leq \min\left\{\frac{b^2\epsilon\epsilon\nabla}{10b\alpha_\epsilon}, \frac{b^2\epsilon\sqrt{\epsilon\nabla}}{4b\alpha_\epsilon\sqrt{p}}, \frac{1}{2}\right\}$, there exist a constant C independent of t such that for all $t \in [0, T_{\max})$ where $T_{\max} < \infty$, we have*

$$(4.20) \quad \left(\int_{M_t} f_\sigma^p d\mu_t \right)^{\frac{1}{p}} \leq C.$$

Proof. For $t \geq t_0$, from (4.10), we have

$$(4.21) \quad \begin{aligned} \frac{\partial}{\partial t} \int_{M_t} f_\sigma^p d\mu_t &\leq \int_{M_t} p f_\sigma^{p-1} \frac{\partial}{\partial t} f_\sigma d\mu_t \\ &\leq -p(p-1) \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t \\ &\quad + 4(1-\sigma)p \int_{M_t} \frac{f_\sigma^{p-1}}{a|H|^2 + \beta_\epsilon c} |H| |\nabla H| |\nabla f_\sigma| d\mu_t \\ &\quad - 2p\epsilon\nabla \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\ &\quad + 2\bar{b}p \int_{M_t} f_\sigma^p d\mu_t + \tilde{b}\sigma p \int_{M_t} |A|^2 f_\sigma^p d\mu_t. \end{aligned}$$

As in (4.16), we have

$$(4.22) \quad \begin{aligned} &4(1-\sigma)p \int_{M_t} \frac{f_\sigma^{p-1}}{a|H|^2 + \beta_\epsilon c} |H| |\nabla H| |\nabla f_\sigma| d\mu_t \\ &\leq \frac{2p}{b\mu} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t + 2p\mu \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t. \end{aligned}$$

Substituting (4.22) to (4.21), letting $\mu = \frac{4}{b(p-1)}$ and $p \geq \max\{2, \frac{8}{b\epsilon\nabla} + 1\}$ we obtain

$$\begin{aligned} \frac{\partial}{\partial t} \int_{M_t} f_\sigma^p d\mu_t &\leq -\frac{p(p-1)}{2} \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t \\ &\quad - p\epsilon\nabla \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\ &\quad + 2\bar{b}p \int_{M_t} f_\sigma^p d\mu_t + \frac{\tilde{b}\sigma\alpha_\epsilon p}{b} \int_{M_t} f_\sigma^p (a|H|^2 + \beta_\epsilon c) d\mu_t. \end{aligned}$$

This together with (4.12) implies

$$(4.23) \quad \begin{aligned} \frac{\partial}{\partial t} \int_{M_t} f_\sigma^p d\mu_t &\leq -p(p-1) \left(\frac{1}{2} - \frac{2\tilde{b}\sigma\alpha_\epsilon}{b^2\eta\varepsilon} \right) \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t \\ &\quad - \left(p\varepsilon\nabla - \frac{(2p\eta+5)\tilde{b}\sigma\alpha_\epsilon p}{b^2\varepsilon} \right) \int_{M_t} \frac{f_\sigma^{p-1}}{(a|H|^2 + \beta_\epsilon c)^{1-\sigma}} |\nabla H|^2 d\mu_t \\ &\quad + 2\bar{b}p \int_{M_t} f_\sigma^p d\mu_t. \end{aligned}$$

Now we pick $\eta = \frac{4\tilde{b}\alpha_\epsilon\sigma}{b^2\varepsilon}$ and let $\sigma \leq \min \left\{ \frac{b^2\varepsilon\varepsilon\nabla}{10b\alpha_\epsilon}, \frac{b^2\varepsilon\sqrt{\varepsilon\nabla}}{4b\alpha_\epsilon\sqrt{p}}, \frac{1}{2} \right\}$. Then (4.23) reduces to

$$\frac{\partial}{\partial t} \int_{M_t} f_\sigma^p d\mu_t \leq 2\bar{b}p \int_{M_t} f_\sigma^p d\mu_t.$$

This implies

$$(4.24) \quad \int_{M_t} f_\sigma^p d\mu_t \leq e^{2\bar{b}pt} \int_{M_{t_0}} f_\sigma^p d\mu_t.$$

If $t \in [0, t_0]$, by the smoothness of the mean curvature flow we see that $\int_{M_t} f_\sigma^p d\mu_t$ is bounded. For $t \geq t_0$, we only have to show that T_{\max} is finite.

Lemma 4.5. *The maximal existence time T_{\max} of the mean curvature flow is finite.*

Proof. Fixed a point $y \in \mathbb{F}^{n+d}(c)$ and let r be the distance function on $\mathbb{F}^{n+d}(c)$ from y . Denote also by r the composition $r \circ F_t$. We may assume that $r > 0$ on M_t for $t \in [0, T_{\max})$. In fact, if $d = 1$, we may choose y such that it is outside of a geodesic ball in $\mathbb{F}^{n+1}(c)$ that encloses M_0 . By the maximum principle we see that y doesn't lie in any M_t . If $d > 1$, then the Hausdorff dimension of $F(M \times [0, T_{\max}))$ is no more than $n+1$. So we can also pick a point y such that it doesn't lie in any M_t . In both cases, we have $r > 0$ on each M_t .

From [4], we know that

$$(4.25) \quad \Delta r = \langle H, \partial_r \rangle + \text{co}_c(r)(n - |\partial_r^T|^2).$$

Here $\text{co}_c(r) = \frac{\sqrt{-c} \cosh(\sqrt{-c}r)}{\sinh(\sqrt{-c}r)} = \frac{\sqrt{-c}(e^{\sqrt{-c}r} + e^{-\sqrt{-c}r})}{e^{\sqrt{-c}r} - e^{-\sqrt{-c}r}}$, ∂_r is the gradient of r in $\mathbb{F}^{n+d}(c)$, and ∂_r^T is the tangent part of ∂_r to M_t . Clearly we have $\text{co}_c(r) \geq \sqrt{-c}$ and $|\partial_r^T|^2 \leq 1$.

On the other hand, since F_t satisfies (1.1), we have

$$(4.26) \quad \frac{\partial}{\partial t} r = \langle H, \partial_r \rangle.$$

Combining (4.25) and (4.26) we obtain

$$(4.27) \quad \frac{\partial}{\partial t} r = \Delta r - \text{co}_c(r)(n - |\partial_r^T|^2).$$

Suppose that $r(0) < R$. By the maximum principle we see that

$$(4.28) \quad r(t) < R - (n-1)\sqrt{-c}t.$$

Then $T_{\max} < \frac{R}{(n-1)\sqrt{-c}}$, i.e., the maximal existence time of the mean curvature flow is finite. \square

By Proposition 4.5, we finish the proof of Lemma 4.4. \square

Now we can proceed as in [11] or [14] via a Stampacchia iteration procedure to complete the proof of Theorem 4.1. \square

5. A GRADIENT ESTIMATE FOR THE MEAN CURVATURE

We establish a gradient estimate for the mean curvature flow, which will be used to compare the mean curvature at different points of the submanifold. We also assume that $c < 0$.

Theorem 5.1. *For every $\eta > 0$, there exists a constant C_η independent of t such that for all $t \in [0, T_{\max})$, there holds*

$$(5.1) \quad |\nabla H|^2 \leq \eta |H|^4 + C_\eta.$$

Proof. By direct computation, we have

$$(5.2) \quad \frac{\partial}{\partial t} |H|^4 \geq \Delta |H|^4 - 12 |H|^2 |\nabla H|^2 + \frac{4}{n} |H|^6 + 4nc |H|^4,$$

$$(5.3) \quad \frac{\partial}{\partial t} |\nabla H|^2 \leq \Delta |\nabla H|^2 + C_1 |H|^2 |\nabla A|^2 + C_2 |\nabla A|^2,$$

for constants C_1 and C_2 independent of t .

We also have the following estimate for sufficiently large positive constants N_1 and N_2 independent of t .

$$(5.4) \quad \begin{aligned} \frac{\partial}{\partial t} \left((N_1 + N_2 |H|^2) |\dot{A}|^2 \right) &\leq \Delta \left((N_1 + N_2 |H|^2) |\dot{A}|^2 \right) - \frac{4(n-1)}{3n} (N_2 - 1) |H|^2 |\nabla A|^2 \\ &\quad - \frac{4(n-1)}{3n} (N_1 - C(N_2)) |\nabla A|^2 \\ &\quad - C_2(N_1, N_2) |\dot{A}|^2 (|H|^4 + 1) - 2nc N_1 |\dot{A}|^2. \end{aligned}$$

In (5.4), $C(N_2)$ and $C(N_1, N_2)$ are constants depending on N_2 and N_1 , N_2 respectively. Consider the function $f = |\nabla H|^2 + (N_1 + N_2 |H|^2) |\dot{A}|^2 - \eta |H|^4$. From (5.2), (5.3) and (5.4), we have

$$(5.5) \quad \begin{aligned} \frac{\partial}{\partial t} f &\leq \Delta f - \frac{4(n-1)}{3n} (N_2 - 1) |H|^2 |\nabla A|^2 - \frac{4(n-1)}{3n} (N_1 - C_3(N_2)) |\nabla A|^2 \\ &\quad + C_4(N_1, N_2) |\dot{A}|^2 (|H|^4 + 1) + 12\eta |H|^2 |\nabla H|^2 - \frac{4\eta}{n} |H|^6 - 4nc\eta |H|^4. \end{aligned}$$

Here we have consumed $C_1 |H|^2 |\nabla A|^2 + C_2 |\nabla A|^2$ by firstly choosing sufficiently large N_2 and secondly choosing sufficiently large N_1 . Notice that $|\nabla H|^2 \leq n |\nabla A|^2$. We can choose larger N_2 and N_1 depending on η to consume $12\eta |H|^2 |\nabla H|^2$ and make the second, third terms of the right hand side of (5.5) negative. Since $|\dot{A}|^2 \leq C_0 |H|^{2-\sigma_0}$ for $t \geq t_0$ and $|\dot{A}|^2$ is uniformly bounded for $t \in [0, t_0]$, using Young's inequality we get

$$C_4(N_1, N_2) |\dot{A}|^2 (|H|^4 + 1) - 4nc\eta |H|^4 \leq \frac{4\eta}{n} |H|^6 + C_\eta.$$

Here C_η is a constant depending on η and other quantities but independent of t . Then we obtain

$$\frac{\partial}{\partial t} f \leq \Delta f + C_5.$$

Notice that T_{\max} is finite. Then the theorem follows from the maximum principle and the definition of f . \square

6. CONVERGENCE OF MCF IN A HYPERBOLIC SPACE

Theorem 6.1. *Let $F : M^n \rightarrow \mathbb{F}^{n+d}(c)$ be a smooth closed submanifold, where $n \geq 2$ and $c < 0$. Assume F satisfies*

$$(6.1) \quad |A|^2 \leq \begin{cases} \frac{4}{3n}|H|^2 + \frac{n}{2}c, & n = 2, 3, \\ \frac{1}{n-1}|H|^2 + 2c, & n \geq 4. \end{cases}$$

Then $F_t(M)$ converges to a round point in finite time.

Proof. By the curvature estimate in [5], we see that

$$K_{\min}(x) \geq \frac{1}{2} \left(\frac{1}{n-1} - \alpha_\epsilon \right) |H|^2(x) + \frac{1}{2} (2 - \beta_\epsilon) c.$$

By our choices of α_ϵ and β_ϵ , and the preserved pinching condition, we see that there exists a positive constant ε_0 independent of t such that

$$(6.2) \quad K_{\min}(x) \geq \varepsilon_0 |H|^2.$$

Since T_{\max} is finite, $\max_{M_t} |A|^2 \rightarrow \infty$ as $t \rightarrow T_{\max}$. By similar arguments as in [1, 11, 12] we have $\frac{\max_{M_t} |H|}{\min_{M_t} |H|} \rightarrow 1$ as $t \rightarrow T_{\max}$, and M_t 's converge to a single point o as $t \rightarrow T_{\max}$. If we take a rescaling around o (since $\mathbb{F}^{n+d}(c)$ can be consider as a linear space that isomorphic to \mathbb{R}^{n+d}) such that the total area of the expanded submanifolds are fixed, then the rescaled immersions converge to a totally umbilical immersion as $t \rightarrow T_{\max}$. \square

When $p = 1$ and $n = 3$, we have the following proposition.

Proposition 6.2. *Let $F : M^3 \rightarrow \mathbb{F}^4(c)$ be a smooth closed hypersurface in a hyperbolic space with constant curvature $c < 0$. Assume F satisfies*

$$|A|^2 \leq \frac{1}{2} |H|^2 + 2c.$$

Then the mean curvature flow with F as initial value converges to a round point in finite time.

Proof. If $d = 1$, then $\mathring{A}_I = 0$. If we take $\alpha \leq \frac{1}{n-1}$ and $\beta \geq 2$ for all $n \geq 2$, the left hand side of (3.2) is nonpositive. Hence $|A|^2 \leq \frac{1}{n-1} |H|^2 + 2c$ is preserved along the mean curvature flow for all $n \geq 2$. When $n = 3$, if we set $a = \frac{1}{3(2+\epsilon)}$, then $\epsilon_\nabla = \frac{3}{4} - \frac{1}{3} - a > 0$, and Theorem 4.1 also holds. Then Theorem 5.1 follows. By a similar argument as in the proof of Theorem 6.1, we get the convergence of the mean curvature flow. \square

Remark 6.3. When $d = 1$ and $n = 3$, the pinching condition $|A|^2 \leq \frac{1}{2}|H|^2 + 2c$ is better than condition (6.1). In fact, we have $\frac{1}{2}|H|^2 + 2c - \frac{4}{9}|H|^2 - \frac{3}{2}c = \frac{1}{18}|H|^2 + \frac{1}{2}c > 0$. Here we have used the fact that $|H|^2 + 9c > 0$, which is implied by $|A|^2 \leq \frac{1}{2}|H|^2 + 2c$.

REFERENCES

- [1] B. Andrews and C. Baker: Mean curvature flow of pinched submanifolds to spheres, *J. Differential Geom.* **85**(2010), 357-395.
- [2] C. Baker: The mean curvature flow of submanifolds of high codimension, arXiv: math.DG/1104.4409.
- [3] K. Brakke: The motion of a surface by its mean curvature, Princeton, New Jersey: Princeton University Press, 1978.
- [4] F. J. Carreras, F. Giménez and V. Miquel: Immersions of compact riemannian manifolds into a ball of a complex space form, *Math. Z.* **225**(1997), 103-113.
- [5] B. Y. Chen: Some pinching and classification theorems for minimal submanifolds, *Arch. Math.* **60**(1993), 568-578.
- [6] Y. G. Chen, Y. Giga and S. Goto: Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Differential Geom.* **33**(1991), 749-786.
- [7] L. C. Evans and J. Spruck: Motion of level sets by mean curvature, I, *J. Differential Geom.* **33**(1991), 635-681.
- [8] J. Erbacher: Reduction of the codimension of an isometric immersion, *J. Differential Geom.* **5**(1971), 333-340.
- [9] H. P. Fu and H. W. Xu: Vanishing and topological sphere theorems for submanifolds in a hyperbolic space, *Intern. J. Math.* **19**(2008), 811-822.
- [10] J. R. Gu and H. W. Xu: The sphere theorems for manifolds with positive scalar curvature, arXiv: math.DG/1102.2424.
- [11] G. Huisken: Flow by mean curvature of convex surfaces into spheres, *J. Differential Geom.* **20**(1984), 237-266.
- [12] G. Huisken: Contracting convex hypersurfaces in Riemannian manifolds by their mean curvature, *Invent. Math.* **84**(1986), 463-480.
- [13] G. Huisken: Deforming hypersurfaces of the sphere by their mean curvature, *Math. Z.* **195**(1987), 205-219.
- [14] G. Huisken and C. Sinestrari: Mean curvature flow singularities for mean convex surfaces, *Calc. Var.* **8**(1999), 1-14.
- [15] B. Lawson: Local rigidity theorems for minimal hypersurfaces, *Ann. Math.* **89**(1969), 179-185.
- [16] K. F. Liu, H. W. Xu, F. Ye and E. T. Zhao: The extension and convergence of mean curvature flow in higher codimension, arXiv: math.DG/1104.0971.
- [17] W. W. Mullins: Two-dimensional motion of idealized grain boundaries, *J. Appl. Phys.* **27**(1956), 900-904.
- [18] K. Shiohama and H. W. Xu, The topological sphere theorem for complete submanifolds, *Compositio Math.* **107**(1997), 221-232.
- [19] K. Smoczyk: Longtime existence of the Lagrangian mean curvature flow, *Calc. Var.* **20**(2004), 25-46.
- [20] K. Smoczyk: Mean curvature flow in higher codimension - Introduction and survey, arXiv: math.DG/1104.3222v2.
- [21] K. Smoczyk and M. T. Wang: Mean curvature flows for Lagrangian submanifolds with convex potentials, *J. Differential Geom.* **62**(2002), 243-257.
- [22] M. T. Wang: Mean curvature flow of surfaces in Einstein four-manifolds, *J. Differential Geom.* **57**(2001), 301-338.
- [23] M. T. Wang: Long-time existence and convergence of graphic mean curvature flow in arbitrary codimension, *Invent. math.* **148**(2002), 525-543.
- [24] M. T. Wang: Lectures on mean curvature flows in higher codimensions, *Handbook of geometric analysis. No. 1*, 525-543, *Adv. Lect. Math. (ALM)*, 7, Int. Press, Somerville, MA, 2008.

- [25] H. W. Xu and J. R. Gu: An optimal differentiable sphere theorem for complete manifolds, Math. Res. Lett. **17**(2010), 1111-1124.
- [26] H. W. Xu, F. Ye and E. T. Zhao: Extend mean curvature flow with finite integral curvature, to appear in Asian J. Math. 2011.
- [27] H. W. Xu, F. Ye and E. T. Zhao: The extension for mean curvature flow with finite integral curvature in Riemannian manifolds, to appear in Sci. China Math. 2011.
- [28] H. W. Xu and E. T. Zhao: Topological and differentiable sphere theorems for complete submanifolds, Comm. Anal. Geom. **17**(2009), 565-585.

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLES
REPUBLIC OF CHINA; DEPARTMENT OF MATHEMATICS, UCLA, Box 951555, LOS ANGELES, CA,
90095-1555

E-mail address: `liu@cms.zju.edu.cn`, `liu@math.ucla.edu`

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLES
REPUBLIC OF CHINA

E-mail address: `xuhw@cms.zju.edu.cn`

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLES
REPUBLIC OF CHINA

E-mail address: `yf@cms.zju.edu.cn`

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLES
REPUBLIC OF CHINA

E-mail address: `zhaoet@cms.zju.edu.cn`